VOL. 691, NOS. 1 + 2 JOURNAL OF CHROMATOGRAPHY A 3 FEBRUARY 1995

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Awareness in Biological Sciences (CABS), Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

18TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, MINNEAPOLIS, MN, USA, 8–13 MAY 1994

Foreword
by L.D. Bowers (Indianapolis, IN, USA) 1

GENERAL, THEORY AND DETECTION

High pH mobile phase effects on silica-based reversed-phase high-performance liquid chromatographic columns
by J.J. Kirkland (Newport, DE, USA) and M.A. van Straten and H.A. Claessens (Eindhoven, Netherlands) . . . 3

Use of chromatographic system peaks for continuous quantitative analysis
by N. Mizrotsky, L. Kristol and E. Grushka (Jerusalem, Israel) 21

Microdialysis sampling coupled on-line to fast microbore liquid chromatography
by A. Chen and C.E. Lunte (Lawrence, KS, USA) 29

Optical properties of axial-illumination flow cells for simultaneous absorbance–fluorescence detection in micro liquid chromatography
by A.A. Abbas and D.C. Shelly (Lubbock, TX, USA) 37

Amperostatic–potentiometric detection for micro high-performance liquid chromatography
by A. Siddiqui and D.C. Shelly (Lubbock, TX, USA) 55

Multivariate visualization in the size-exclusion chromatography and pattern recognition of biological samples
by R.D. Ricker and L.A. Sandoval (Newport, DE, USA), J.D. Justice (Mountain View, CA, USA) and F.O. Geiser (Glen Mills, PA, USA) 67

SORBENTS, COLUMNS, PROCEDURES CHIRAL SEPARATIONS

Characterization of a new reversed-phase chromatographic column on a 2-μm porous microspherical silica gel
by H. Moriyama, M. Anegayama, K. Komiya and Y. Kato (Yamaguchi, Japan) 81

Nano-scale design of novel stationary phases to enhance selectivity for molecular shape and size in liquid chromatography
by K. Jinno, K. Nakagawa, Y. Saito, H. Ohta, H. Nagashima and K. Itoh (Toyohashi, Japan) and J. Archer and Y.-L. Chen (Folsom, CA, USA) 91

Displacement chromatography in biotechnological downstream processing
by R. Freitag and J. Breier (Hannover, Germany) 101

Preparation and use of immunoglobulin-binding affinity supports on Emphaze beads
by G.T. Hermanson, G.R. Mattson and R.I. Krohn (Rockford, IL, USA) 113

Effect of resin sulfonation on the retention of polar organic compounds in solid-phase extraction
by P.J. Dumont and J.S. Fritz (Ames, IA, USA) 123

Methods and materials for solid-phase extraction
by J.S. Fritz, P.J. Dumont and L.W. Schmidt (Ames, IA, USA) 133

Characteristics and applications of a new high-performance liquid chromatography guard column
by M. Capparella, W. Foster III, M. Larrousse, D.J. Phillips, A. Pomfret and Y. Tuvim (Milford, MA, USA) . . 141

Reversed-phase high-performance liquid chromatography using enhanced-fluidity mobile phases
by Y. Cui and S.V. Olesik (Columbus, OH, USA) 151

Modification of a conventional high-performance liquid chromatography autoinjector for use with capillary liquid chromatography (Short communication)
by R.C. Simpson (King of Prussia, PA, USA) 163

Solvent modulation in liquid chromatography: extension to serially coupled columns
by P.H. Lukulay and V.L. McGuffin (East Lansing, MI, USA) 171

Chiral separation retention mechanisms in high-performance liquid chromatography using bare silica stationary phase and β-cyclodextrin as a mobile phase additive
by R.H. Pullen (Atlanta and Marietta, GA, USA), J.J. Brennan (Marietta, GA, USA) and G. Patonay (Atlanta, GA, USA) 187

Laser-based dynamic surface tension detection for liquid chromatography by probing a repeating drop radius
by L.R. Lima III and R.E. Synovec (Seattle, WA, USA) 195

Chemometric characterization of Lewis base-modified zirconia for normal phase chromatography
by D.A. Whitman, T.P. Weber and J.A. Blackwell (St. Paul, MN, USA) 205

High-performance chiral displacement chromatographic separations in the normal-phase mode. III. Separation of the enantiomers of 5-vinylpyrrolidin-2-one using the Chiralcel-OD stationary phase
by P.L. Camacho-Torralba and Gy. Vigh (College Station, TX, USA) 213

CARBOXYLIC ACIDS, PROSTAGLANDINS, LEUKOTRIENES

Comparison of the retention of organic acids on alkyl and alkylamide chemically bonded phases
by T. Czajkowska and I. Hrabovsky (Princeton, NJ, USA) and B. Buszewski, R.K. Gilpin and M. Jaroniec (Kent, OH, USA) 217

Use of secondary chemical equilibria in liquid chromatography to determine dissociation constants of leukotriene B_4 and prostaglandin B_2
by J.E. Hardcastle, M. He, B. Begum and R. Vermillion-Salsbury (Denton, TX, USA) 225

Simultaneous sorption and analytical derivatization on a polystyrene–divinylbenzene polymer. Preparation of chromophoric and fluorophoric derivatives of the prostaglandins
by J.M. Rosenfeld and X. Fang (Hamilton, Canada) 231

STEROIDS

Use of particle-loaded membranes to extract steroids for high-performance liquid chromatographic analyses. Improved analyte stability and detection
by G.L. Lensmeyer, C. Onsager, I.H. Carlson and D.A. Wiebe (Madison, WI, USA) 239

High-performance liquid chromatographic analysis of human erythrocyte oxysterols as Δ^4-3-ketone derivatives
by J.I. Teng and L.L. Smith (Galveston, TX, USA) 247

PEPTIDES, PROTEINS, ENZYMES

High-performance liquid chromatographic determination of sulfated peptides in human hemofiltrate using a radioactivity monitor
by A.G. Schepky, P. Schulz-Knappe and W.-G. Forssmann (Hannover, Germany) 255

High-performance liquid chromatography of amino acids, peptides and proteins. CXXXVIII. Adsorption of horse heart cytochrome *c* onto a tentacle-type cation exchanger
by J. Xie, M.-I. Aguilar and M.T.W. Hearn (Clayton, Australia) 263

High-performance liquid chromatography of amino acids, peptides and proteins. CXXXIX. Impact of operating parameters in large-scale chromatography of proteins
by Q.-M. Mao, I.G. Prince and M.T.W. Hearn (Clayton, Australia) 273

Compositional analysis of the phenylthiocarbamyl amino acids by liquid chromatography–atmospheric pressure ionization mass spectrometry with particular attention to the cyst(e)ine derivatives
by K. Schmeer (Tübingen, Germany), M. Khalifa, J. Császár and G. Farkas (Budapest, Hungary), E. Bayer (Tübingen, Germany) and I. Molnár-Perl (Budapest, Hungary) 285

Sensitivity and selectivity of the electrochemical detection of the copper(II) complexes of bioactive peptides, and comparison to model studies by rotating ring-disc electrode
by J.-G. Chen, S.J. Woltman and S.G. Weber (Pittsburgh, PA, USA) 301

Separation of amino acids, peptides and proteins on molecularly imprinted stationary phases
by M. Kempe and K. Mosbach (Lund, Sweden) 317

Separation and analysis of proteins by perfusion liquid chromatography and electrospray ionization mass spectrometry
by J.F. Banks, Jr. (Branford, CT, USA) 325

High-performance liquid chromatographic assay of glycosyltransferases using flavonoids as substrate
by M. Pace, D. Agnellini, C. Gardana, P.L. Mauri and P.G. Pietta (Milan, Italy) 331

Selectivity optimization of reversed-phase high-performance liquid chromatographic peptide and protein separations by varying bonded-phase functionality
by B.E. Boyes (Newport, DE, USA) and D.G. Walker (Vancouver, Canada) 337

Insights into the role of the hydrogen bond and hydrophobic effect on recognition in molecularly imprinted polymer synthetic peptide receptor mimics
by I.A. Nicholls, O. Ramström and K. Mosbach (Lund, Sweden) 349